



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

10/526 849
Rec'd PCT/PTO 04 MAR 2005
PCT/NL 03/00622

#2

REC'D 30 SEP 2003

WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02078645.5

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 02078645.5
Demande no:

Anmeldetag:
Date of filing: 05.09.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Pantarhei Bioscience B.V.
P.O. Box 464
3700 AL Zeist
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Method of treating neurological and mental disorders

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

METHOD OF TREATING NEUROLOGICAL AND MENTAL DISORDERS

EPO - DG 1

05.09.2002

FIELD OF THE INVENTION

(95)

5

The present invention is concerned with a method of treating neurological and/or mental disorders that are associated with, or pathogenetically related to deficient serotonin neurotransmission. More particularly the invention relates to a method of increasing cerebral intercellular serotonin levels and/or of preventing decreases in such serotonin levels by a method that comprises administering a serotonin re-uptake inhibitor.

10

BACKGROUND OF THE INVENTION

15

Serotonin (5-hydroxytryptamine) is a neurotransmitter that is found in relatively high concentrations in the lateral gray horns of the spinal cord and in a number of areas in the brain. A system of serotonin-containing neurons that have their cell bodies in the raphe nuclei of the brain stems project to portions of the hypothalamus, the limbic system, the neocortex and the spinal cord. It has been demonstrated that serotonin interacts with a great number of receptors in the brain and controls or affects processes which regulate many bodily organs and functions.

20

Serotonin is an important chemical messenger participating in the transmission of nerve impulses in the brain. These messengers are released at specific sites on pre-synaptic cells and received, to complete transmission of the impulse, at specific sites on post-synaptic cells. Their effect is then terminated by metabolism or by uptake into the pre-synaptic cells. In addition, some of the released serotonin is inactivated by monoamine oxidase to form 5-hydroxyindoleacetic acid. This is the principal urinary metabolite of serotonin.

25

Serotonin is formed in the body by hydroxylation and decarboxylation of the essential amino acid L-tryptophan. In the biosynthesis of serotonin from L-tryptophan, L-tryptophan is hydroxylated in the presence of the enzyme tryptophan hydroxylase to form the intermediate product L-5-hydroxytryptophan. This intermediate product is decarboxylated in the presence of the enzyme 5-hydroxytryptophan decarboxylase to form serotonin.

30

Serotonin deficiencies in the brain have been associated with a number of mainly

psychological disorders such as (endogenous) depression, insomnia, excessive appetite and lowered pain threshold.

An important discovery in medicinal chemistry of the past decades are the so called serotonin re-uptake inhibitors (SRI's), which are particularly effective in the treatment of depression. SRI's increase the availability of serotonin in the synapse by reducing the uptake
5 of serotonin by the serotonin uptake carrier (transporter protein). Dysfunction of the serotonin - neurons resulting from excessive uptake results in depression, as well as other pathologies of the central nervous system.

Among the commercially available SRI's are fluoxetine, nefazodone, sertraline,
10 venlafaxine, citalopram, fluvoxamine, paroxetine and trazodone. While the primary activity of these drugs is the inhibition of the re-uptake of serotonin, the cascade of monoamine processes in the brain connects serotonin with both norepinephrine and dopamine. Thus, the increase of availability of serotonin may result in increased availability of norepinephrine and dopamine as well.

15 Since the commercial introduction of the first SRI's in the early eighties these pharmaceutical components have found widespread application, particularly as anti-depressants. Despite this success there are a few serious drawbacks to the use of SRI's that have not yet been resolved. One such drawback relates to the fact that after administration of SRI has commenced, it usually takes a few weeks before its therapeutic impact (action)
20 becomes apparent. Consequently, these drugs are less suited for treating individuals who suffer from non-chronic afflictions or for treating individuals who need urgent relief (e.g. suicidal patients).

Another drawback of known SRI's is related to the relatively low responder rate that has been observed for these drugs. In general it is found that only about 60-70% of the
25 patients treated with SRI respond well to such treatment. Given that it takes a few weeks before it can be established whether or not a patient responds well to treatment with SRI, it will be clear that better responder rates are highly desirable.

In view of the aforementioned drawbacks there is a need for a method of treatment that will deliver the functionality of SRI's more quickly and/or with a higher responder rate. It is
30 an objective of the present invention to make available a method that meets these requirements.

SUMMARY OF THE INVENTION

The inventors have surprisingly found that the aforementioned objective can be realised by a method of treatment that comprises the administration of an effective amount of a combination of SRI and a vitamin B6 component. The co-administration of said vitamin B6 component(s) was unexpectedly found to advance the onset of action of the SRI. Also it appears that the co-administration of the vitamin B6 component improves the responder rate of a given SRI.

The present method can advantageously be used for the treatment of any disorders or diseases for which SRI's have been applied successfully. Such disorders and diseases include (endogenous) depression, insomnia, excessive appetite, lowered pain threshold etc.

Although the combined use of SRI's and vitamin B6 has been described in the prior art, it has not been recognised before that vitamin B6 has the capability of advancing the onset of the action of SRI's or of increasing the responder rate of these drugs.

US 4,596,807 (Crosby) describes a method for controlling pain, depression and sedation, which method comprises administering a composition comprising a serotonin precursor, such as L-tryptophan or L-5-hydroxytryptophan in an amount effective to increase the brain serotonin to a supra normal level, in combination with a SRI in an amount effective to inhibit the re-uptake of serotonin. The composition may additionally comprise vitamin B6 (pyridoxine) and vitamin C (ascorbic acid) so as to aid biosynthesis of serotonin in case of patients suffering from vitamin depletion. In accordance with this patent administration of the composition is desirably effected in 1-4 portions daily, delivering 100-800 mg per day of SRI and 40-400 mg per day of the pyridoxine.

US 5,885,976 (Sandyk) is concerned with methods for the treatment of neurological and mental disorders related to deficient serotonin neurotransmission and impaired pineal melatonin functions. The method described comprises administering an effective amount of a composition which increases serotonin transmission followed by the application to the brain of a sufficient amount of an AC pulsed magnetic field to treat the disorder. It is mentioned that the composition may contain a stimulant of serotonin synthesis which is vitamin B1, vitamin B3, vitamin B6, biotin, S-adenosyl methionine, vitamin D, folic acid, ascorbic acid, magnesium, coenzyme Q10, piracetam, or mixtures of two or more thereof. Furthermore it is stated that the composition can include a SRI which is sertraline, nefazodone, trazodone, fluoxetine or a mixture thereof. The ranges of daily dosage levels specifically mentioned for a

number of SRI's in this patent broadly cover the range of 25-600 mg.

The above prior art publications that describe the combined use of SRI's and vitamin B6 teach to employ SRI's at relatively high daily dosage levels of at least 100 mg. Presumably these high dosages are employed to push forward the onset of action and/or to achieve maximum responder rate. The method according to the present invention does not require such high doses to achieve the same, due to the co-administration with the vitamin B6 component nor does it require the application of an AC pulsed magnetic field as taught by US 5,885,976 (Sandyk).

DETAILED DESCRIPTION OF THE INVENTION

Accordingly, one embodiment of the present invention relates to a pharmaceutical formulation for use in a method of increasing cerebral intercellular serotonin levels and/or of preventing decreases in such serotonin levels, said method comprising the administration of:

- serotonin re-uptake inhibitor in a daily amount of between 0.4 and 80 mg and
- vitamin B6 component in a daily amount of between 0.01 and 10 mmoles,

wherein the method does not include the application to the brain of an AC pulsed magnetic field of at least 7.5 picotesia flux density for at least 15 minutes. More preferably the method does not include the application of an AC pulsed magnetic field of at least 7.5 picotesia. Most preferably the method does not include the application of an AC pulsed magnetic field at all.

The term "serotonin re-uptake inhibitor", also sometimes referred to as "serotonin selective re-uptake inhibitor (or SSRI)" encompasses those re-uptake inhibitors that are capable of significantly inhibiting re-uptake of serotonin by blocking its transporter protein, by inhibiting monoamine oxidase and/or by (selectively) blocking cerebral serotonin receptors. Preferably the SRI used in the present method is capable of blocking the serotonin transporter protein.

The term "on a daily basis", when used in connection with a mentioned dosage amount, should not be interpreted restrictedly. For instance, the above mentioned requirement that the administration of the present formulation is to provide, on a daily basis, less than 80 mg SRI, encompasses a protocol wherein SRI is administered once a week, provided the weekly dosage is less than 560 mg, i.e. such that the average daily dose is less than 80 mg SRI.

The term "vitamin B6 component" as used throughout this document encompasses any components which *in vivo*, particularly once these components or their metabolites have entered the bloodstream, are converted into pyridoxal or a pyridoxal salt. Particularly useful are vitamin B6 components that *in vivo* are converted for at least 10 mol% into pyridoxal or a pyridoxal salt within 24 hours after administration.

Inside living human and animal cells, pyridoxal phosphate and pyridoxamine phosphate are the biologically active forms of vitamin B6, acting as a co-enzyme in more than 100 biological reactions. In the form of pyridoxal 5'-phosphate and pyridoxamine 5'-phosphate, vitamin B6 acts as the coenzyme of a series of enzymes, which catalyse transamination, decarboxylation, deamination, desulphydration and the cleavage or synthesis of amino acids. The aminotransferases represent an important link between amino acid, carbohydrate and fatty acid metabolism and the energy-producing citric acid cycle. The decarboxylases convert amino acids to the corresponding biogenic amines, such as histamine, hydroxytyramine, serotonin, γ -aminobutyric acid, ethanolamine and taurine, some of which are substances of high physiological activity [regulation of the blood vessel diameter, neurohormonal actions, essential components of phospholipids and bile acids]. In accord with the general importance these enzymatic reactions take place in virtually all organs most intensively in the liver, heart and brain. (Lit. Vitamin Compendium; Roche, Vitamins and Chemicals Department. F. Hoffmann- La Roche and Co. Ltd, Basle, Switzerland).

As described herein before SRI and the vitamin B6 component may be co-administered in a sequential as well as in a coextensive fashion. Hence the term "pharmaceutical formulation" as used throughout this document should be construed to encompass pharmaceutical kits which comprise separate dosage units for the SRI component and the vitamin B6 component. Preferably, however, the SRI component and vitamin B6 component are combined into a single dosage unit.

The SRI and vitamin B6 component may suitably be administered enterally (e.g. orally or rectally), parenterally (e.g. intravenous or intramuscular) or topically. Preferably, the present method employs enteral administration, most preferably oral administration. The present method may suitably be used to treat a wide variety of mammals. The method, however, is particularly suitable for treating humans.

The present method is particularly useful for treating depression, anxiety disorders (e.g. generalised anxiety disorder (GAD), panic disorder and social anxiety disorder (social

phobia)), obsessive compulsive disorder, premenstrual syndrome, post traumatic stress disorder, pain, sedation, insomnia, weight gain and excessive appetite. Preferably the present method is used for the treatment of depression, anxiety disorders, panic disorder, or obsessive compulsive disorder. Most preferably the present method is used for the treatment of
5 depression. The present method does not encompass the treatment of hot flushes in e.g. hypo-estrogenic females and androgen-deprived males.

Depression is often associated with other diseases and conditions, or caused by such other conditions. Depression may also be associated with abuse of any substance, or may be associated with behavioural problems resulting from or occurring in combination with head
10 injuries, mental retardation or stroke. Depression in all its variations is a target of treatment for formulations according to the present invention.

According to another preferred embodiment of the method of treatment according to the invention said method comprises at least once daily administration of the pharmaceutical formulation. Since SRI's are metabolised relatively slowly, it is possible to apply longer
15 intervals between doses. However, longer intervals will usually lead to an increase in amplitude of the observed fluctuations of SRI blood serum levels. Since a minimum blood serum level needs to be maintained to achieve the desired results, the maximally occurring blood serum level of SRI will inevitably increase with incrementing dosage intervals. This is not in line with the aim of the present invention to not employ unnecessarily high SRI blood
20 serum levels.

The present method of increasing cerebral serotonin levels offers the advantage that it employs relatively low levels of SRI. Thus in comparison to ordinary methods, employing higher dosage of SRI, the present method exhibits a decreased incidence of side-effects. This may be explained from the fact that the vitamin B6 component, in the proposed dose ranges,
25 has no side-effects. In a preferred embodiment of the invention the method provides on a daily basis a minimum amount of 0.6 mg SRI. The preferred daily maximum amount is 70 mg SRI. More preferably the daily maximum amount is 60 mg SRI, even more preferably it is 50 mg. Most preferably the maximum daily amount of SRI provided by the method is 40 mg. The daily minimum amount provided by the present method preferably is 1 mg, more preferably 2
30 mg and most preferably 3 mg.

According to another preferred embodiment the present method provides on a daily basis a minimum amount of 3.0 mg SRI, calculated as trazodone equivalent. More preferably the daily minimum amount is 10.0 mg trazodone equivalent, most preferably 15.0 mg

trazodone equivalent. The preferred daily maximum amount is equivalent to less than 90 mg trazodone, more preferably less than 80 mg trazodone. Most preferably the maximum daily amount of SRI provided by the method is equivalent to less than 75 mg trazodone.

In order to facilitate the translation of given amounts of trazodone into the equivalent amounts of another SRI and vice versa, the following table provides the conversion factors that are to be used:

	Trazodone conversion factor
citalopram	0.15
fluoxetine	0.15
fluvoxamine	0.7
paroxetine	0.15
sertraline	0.3
venlafaxine	0.5
trazodone	1.0
mirtazapine	0.2
nefazodone	1.3

With the help of the above table it can be calculated that 100 mg trazodone is equivalent to 30 mg sertraline or 15 mg fluoxetine.

The present formulations preferably do not contain a narcotic, such as those mentioned in US 4,596,807 (Crosby). More particularly the present method does not encompass the administration of a narcotic selected from the group consisting of codeine, oxycodone, propoxyphene, pentazocine, morphine, meperidine, levorphanol, menthadone and mixtures thereof, in an amount effective to produce analgesia.

The present method of treatment preferably comprises administration of SRI in a daily amount of between 0.01 and 1 mg SRI per kg bodyweight and vitamin B6 component in a daily amount of between 0.001 and 0.2 mmoles per kg bodyweight. More preferably the SRI is administered in a daily amount of between 0.02 and 0.8 mg SRI/kg and the vitamin B6 component in a daily amount of between 0.004 and 0.1 mmoles/kg. Most preferably the SRI is administered in a daily amount of between 0.03 and 0.7 mg SRI/kg.

Analogues of vitamin B6 which can suitably be used in accordance with the present invention are those selected from the group consisting of pyridoxal, pyridoxamine, acetals of pyridoxal, condensation products arising from the reaction of the aldehyde group of pyridoxal

with an amine, and addition salts of any of the foregoing members of the group with pharmaceutically acceptable salts. The term "pharmaceutically acceptable salts" includes salts with pharmaceutically acceptable acids or bases, e.g. acids such as sulphuric, hydrochloric, nitric, phosphoric acid, etc. or bases such as alkali or alkaline earth metal hydroxides, ammonium hydroxides, alkyl ammonium hydroxides etc.

In order to obtain a significant beneficial effect from the co-administration of SRI and vitamin B6 component, it is advisable to apply the vitamin B6 component and SRI in a dosage ratio of at least 0.001 mmole/mg, preferably of at least 0.01 mmole/mg, more preferably of at least 0.03 mmole/mg and most preferably of at least 0.05 mmole/mg.

Generally the aforementioned ratio will not exceed 2 mmole/mg, preferably it will not exceed 1 mmole/mg and more preferably it will not exceed 0.5 mmole/mg.

In a preferred embodiment of the present invention the SRI's employed are selected from the group selected of citalopram, fluoxetine, norfluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine, zimelidine, femoxetine, trazodone, nefazodone, mirtazapine, pharmaceutically acceptable salts of these inhibitors and mixtures thereof. Most preferably the SRI's are selected from the group consisting of citalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine, zimelidine, trazodone, nefazodone, mirtazapine, pharmaceutically acceptable salts of these inhibitors and mixtures thereof. It is noted that the SRI's used in present formulations may be present in the form of a racemic mixture as well as in the form of a stereo-enantiomer.

In another preferred embodiment of the present invention the pharmaceutical formulation additionally comprises at least 300 mg, more preferably from 500 to 10,000 mg of a serotonin precursor selected from the group consisting of L-tryptophan, 5-hydroxytryptophan, precursors of these tryptophan substances and mixtures thereof. The inclusion of the latter precursors may assist in alleviating serotonin deficiency since serotonin is biosynthesised from tryptophane through the following metabolic chain:

tryptophan > 5-hydroxytryptophan > 5-hydroxytryptamine > serotonin.

Tryptophan is usually not transported in the blood in a free state, but rather bound to or complexed with blood serum albumin. Tryptophan is the only circulating amino acid that is significantly bound to human blood serum albumin. It has been shown that salicylates displace tryptophan from its protein binding site on albumin in blood plasma thereby raising the free, circulating tryptophan concentration in blood. This free or unbound tryptophan is more easily converted to serotonin than the bound form. Hence the present pharmaceutical

formulation may advantageously contain as an additional component a salicylate. Preferably the formulation comprises at least 0.05 mmoles of such a salicylate, more preferably from 0.1 to 1.5 mmoles of the salicylate. Here the term salicylate includes both the acid and the salt. The salicylate is preferably selected from the group consisting of sodium salicylate, choline salicylate, magnesium salicylate and mixtures thereof.

Yet another component which may additionally be included in the present formulation - is a component that acts as a stimulant of the serotonin receptor, such as buspirone. Hence in another preferred embodiment the formulation comprises at least 10 μ moles, preferably from 20 to 300 μ moles of buspirone.

The pharmaceutical formulations according to the invention can be solid or semi-solid dosage forms such as tablets, capsules, cachets, pellets, pills, suppositories, powders and granules, as well as fluid dosage forms such as solutions, emulsions, suspensions, ointments, pastes, creams, gels, jellies and foams. In addition to the pharmacologically active components, the formulation according to the invention contains pharmaceutically acceptable excipient, usually in an amount of between 50 and 99.9 wt.%.

Tablets and equivalent solid and semi-solid dosage forms can suitably contain excipients such binders (e.g. hydroxypropylmethyl cellulose, polyvinyl-pyrrolidone, other cellulosic materials and starch), diluents (e.g. lactose and other sugars, starch, dicalcium phosphate and cellulosic materials), disintegrating agents (e.g. starch polymers and cellulosic materials) and lubricating agents (e.g., stearates and talc).

Transdermal delivery systems include patches, gels, tapes and creams, and can contain excipients such as solubilisers, permeation enhancers (e.g. fatty acids, fatty acid esters, fatty alcohols and amino acids), hydrophilic polymers (e.g. polycarbophil and polyvinyl pyrrolidone) and adhesives and tackifiers (e.g. polyisobutylenes, silicone-based adhesives, acrylates and polybutene).

Transmucosal delivery systems include patches, tablets, suppositories, pessaries, gels, and creams, and can contain excipients such as solubilizers and enhancers (e.g. propylene glycol, bile salts and amino acids), and other vehicles (e.g. polyethylene glycol, fatty acid esters and derivatives, and hydrophilic polymers such as hydroxypropylmethyl cellulose and hyaluronic acid).

Injectable delivery systems include solutions, suspensions, gels, microspheres and polymeric injectables, and can comprise excipients such as solubility-altering agents (e.g. ethanol, propylene glycol and sucrose) and polymers (e.g. polycaprylactones, and PLGA's).

Implantable systems include rods and discs, and can contain excipients such as PLGA and polycapryl lactone.

Other delivery systems that can be used for administering the pharmaceutical composition of the invention include intranasal delivery systems such as sprays and powders, and sublingual delivery systems.

The inventors have observed that in order to obtain the full benefit from the co-administration of the vitamin B6 component it is advantageous to administer said component at relatively short time intervals, e.g. every 4-6 hours. It is believed that, following absorption, vitamin B6 is quickly converted into metabolites that have less of a synergistic interaction with the SRI. Consequently, it is advantageous to administer the vitamin B6 component by means of a sustained release formulation, so as to avoid the burden of having to administer said component 4-6 times per day. As a result of the controlled release from a sustained release formulation (e.g. an oral dosage form, a suppository or a transdermal patch) the blood serum level of the vitamin B6 component can be maintained at a stable level which coincided with the level that produces the optimum synergistic interaction between the SRI and the vitamin B6 component.

The present invention encompasses an embodiment wherein the vitamin B6 component is separately administered from the SRI in the form of a slow release formulation. Preferably, however, both active agents are administered in the form of a slow release formulation. Most preferably both agents are administered through a single slow release formulation, in particular a single solid dosage form.

Generally the present sustained release formulation should release the vitamin B6 component at a rate of at least 0.001 mmole/hour during the first 4 hours after administration. Preferably the release rate during said period is at least 0.005 mmole/hour, more preferably at least 0.01 mmole/hour.

In a particularly preferred embodiment the present method comprises the administration of a sustained release formulation at intervals of at least one day. More preferably, said intervals are within the range of 1 day and 1 week. Most preferably the sustained release formulation is administered once daily.

Another embodiment of the present invention relates to a pharmaceutical formulation for oral, rectal, buccal or transdermal administration, comprising a solid dosage form that contains as active agents at least 0.4 mg of serotonin re-uptake inhibitor and at least 0.01 mmole of vitamin B6 component as well as a pharmaceutically acceptable carrier, optionally

in combination with a membrane coating material, that provides for sustained release of these active agents from the dosage form. The term "solid dosage form" also encompasses dosage units that e.g. comprise a liquid or a paste which is encased in a capsule.

5 The pharmaceutical formulation according to the present invention may suitably take the form of an oral dosage unit, a buccal dosage unit, a suppository, a transdermal patch or a depot injection. Preferably said formulation comprises an oral dosage unit or a suppository. Most preferably the present formulation comprises an oral dosage unit.

10 In order to ensure the sustained release of the active agents, the present formulation may advantageously comprise a membrane coating material that separates the active agents from the exterior. In case the formulation is administered orally, buccally or rectally, the active agents will gradually diffuse through said membrane coating and/or said coating will gradually degrade allowing release of the active agents from the interior of the formulation.

15 In a particularly preferred embodiment, the solid dosage form is effective to provide a sustained release profile wherein less than 50 wt.% of the vitamin B6 component is released from the dosage form within the first 4 hours after administration and more than 80 wt.% of the vitamin B6 component is released from the dosage form within 24 hours.

20 Yet another aspect of the present invention relates to a pharmaceutical formulation containing at least 0.4 mg of SRI, at least 0.01 mmole of vitamin B6 component and a pharmaceutically acceptable carrier, wherein vitamin B6 component and SRI are present in a ratio of between 0.01 and 1 mmole/mg, preferably in a ratio of between 0.05 and 0.5 mmole/mg. The application of a relatively high ratio of vitamin B6 component to SRI ensures that optimum responder rates are achieved at a given SRI dosage.

The invention is further illustrated by means of the following examples.

25 EXAMPLE

Example I: Onset of action

30 *Patient Selection*

Patients included in this study have been diagnosed as having Major Depressive Disorder using criteria as set forth in the DSM-IV. Two psychiatrists have confirmed this diagnosis independently before inclusion. All patients are between the ages of 18 and 75

years, are not vitamin B6 deficient and have no major medical problems nor are they using any medication to treat their depression.

The severity of the depression of the patients is assessed at baseline using the Hamilton Rating Scale for Depression (HAM-D 17 items) (for review: Hamilton M. (1960) "A rating scale for depression". J. Neurol. Neurosurg. Psychiatry 23, 56; Hamilton M. (1967) "Development of a rating scale for primary depressive illness". Br J Soc Clin Psychol 6, 278-296). Only patients with a score greater than 21 on the HAM-D 17 continue in the study (a HAM-D 17 score of 18-25 indicates moderate to severe depression). Based on these criteria, twenty (20) patients with a depressive episode are selected for this study.

Dose regimen

The patients are randomised to treatment groups A or B. Each group receives a different daily dose regimen, i.e.:

- A: 20 mg fluoxetine + 500 mg vitamin B6; and

- B: 20 mg fluoxetine + placebo (appearance of the placebo is identical to the tablet containing vitamin B6)

Scoring

During the study period of 10 weeks, patients are assessed using the HAM-D at weekly intervals by experienced raters.

Results

On average, the HAM-D scores of patients in treatment group A are found to decline faster than in treatment group B. This indicates that the regimen of group A has a positive effect on the onset of action of fluoxetine.

Example II: Non-responders

Patient Selection

Patients included in this study were selected on the basis of the same criteria as described in example I.

After 6 weeks of adequate SSRI therapy (i.e., fluoxetine 20 mg daily), those patients

who showed insufficient improvement on the HAM-D 17, are considered to be 'non-responders'. Twenty (20) 'non responders' are selected for inclusion in the present study.

Dose regimen

5 The patients are randomised to treatment groups A or B. Each group receives a different daily dose regimen, i.e.:

- A: 20mg fluoxetine + 500mg vitamin B6; and
- B: 20mg fluoxetine + placebo

10 *Scoring*

During the study period of 8 weeks, patients are assessed using the HAM-D 17.

Results

15 The number of patients showing a decrease of baseline HAM-D scores of 50% or greater in group A is higher than that in group B. This indicates that the regimen of group A enhances the response to fluoxetine treatment in patients considered to be non-responders.

Example III: Onset of action (low dose of SSRI)

20 *Dose regimen*

The patients are randomised to treatment groups A or B. Each group receives a different daily dose regimen, i.e.:

- A: 10mg fluoxetine + 500mg vitamin B6; and
- B: 10mg fluoxetine + placebo

25

Scoring

During the study period of 10 weeks, patients are assessed using the HAM-D 17.

Results

30 The patients in group A are found to show a faster decline in the HAM-D scores than those in treatment group B. This indicates that the regimen of group A has a positive effect on the onset of action of low dosed fluoxetine.

CLAIMS

05.09.2002

(95)

1. A pharmaceutical formulation for use in a method of increasing cerebral intercellular serotonin levels and/or of preventing decreases in such serotonin levels, said method comprising the administration of:
- 5 a. serotonin re-uptake inhibitor in a daily amount of between 0.4 and 80 mg and
b. vitamin B6 component in a daily amount of between 0.01 and 10 mmoles, wherein the method does not include the application to the brain of an AC pulsed magnetic field of at least 7.5 picotesia flux density for at least 15 minutes.
- 10 2. A pharmaceutical formulation according to claim 1 for use in a method of treating depression, anxiety disorders, obsessive compulsive disorder, premenstrual syndrome, post traumatic stress disorder, pain, sedation, insomnia, weight gain and excessive appetite.
- 15 3. A pharmaceutical formulation according to claim 1 or 2, wherein the method comprises at least once daily administration of the SRI and the vitamin B6 component.
- 20 4. A pharmaceutical formulation according to any one of claims 1-3, wherein the method comprises administration of SRI in a daily amount of between 0.01 and 1 mg per kg bodyweight and vitamin B6 component in a daily amount of between 0.001 and 0.2 mmoles per kg bodyweight.
- 25 5. A pharmaceutical formulation according to any one of claims 1-4, wherein the vitamin B6 analogue is selected from the group consisting of pyridoxal, pyridoxamine, acetals of pyridoxal, condensation products arising from the reaction of the aldehyde group of pyridoxal with an amine, and addition salts of any of the foregoing members of the group with pharmaceutically acceptable salts.
- 30 6. A pharmaceutical formulation according to any one of claims 1-5, wherein the dosage ratio of vitamin B6 component to serotonin re-uptake inhibitor is in the range of 0.01 to 1 mmole/mg, preferably in the range of 0.05 to 0.5 mmole/mg.

- 5 7. A pharmaceutical formulation according to any one of claims 1-6, wherein the serotonin re-uptake inhibitor is selected from the group selected of citalopram, fluoxetine, norfluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine, zimelidine, femoxetine, trazodone, nefazodone, pharmaceutically acceptable salts of these inhibitors and mixtures thereof.
- 10 8. A pharmaceutical formulation according to any one of claims 1-7, wherein the formulation comprises at least 300 mg of a serotonin precursor selected from the group consisting of L-tryptophan, 5-hydroxytryptophan, precursors thereof and mixtures thereof.
9. A pharmaceutical formulation according to any one of claims 1-8, wherein the formulation comprises at least 0.05 mmoles of a salicylate.
- 15 10. A pharmaceutical formulation according to any one of claims 1-9, wherein the formulation does not contain a narcotic selected from the group consisting of codeine, oxycodone, propoxyphene, pentazocine, morphine, meperidine, levorphanol, menthadone and mixtures thereof, in an amount effective to produce analgesia.
- 20 11. A pharmaceutical formulation according to any one of claims 1-10, wherein the method comprises the at least once daily oral administration of a slow release formulation containing the vitamin B6 component.
- 25 12. A pharmaceutical formulation for oral, rectal, buccal or transdermal administration, comprising a solid dosage form that contains as active agents at least 0.4 mg of serotonin re-uptake inhibitor and at least 0.01 mmole of vitamin B6 component as well as a pharmaceutically acceptable carrier, optionally in combination with a membrane coating material, that provides for sustained release of these active agents from the dosage form.
- 30 13. A pharmaceutical formulation according to claim 12, wherein the solid dosage form is effective to provide a sustained release profile wherein less than 50 wt.% of the vitamin B6 component is released from the dosage form within the first 4 hours after administration and more than 80 wt.% of the vitamin B6 component is released from the dosage form within 24 hours.

14. A pharmaceutical formulation according to claim 12 or 13, wherein the solid dosage form release the vitamin B6 component at a rate of at least 0.001 mmole/hour during the first 4 hours after administration.

5

15. A pharmaceutical formulation containing at least 0.4 mg of serotonin re-uptake inhibitor, - at least 0.01 mmole of vitamin B6 component and a pharmaceutically acceptable carrier, wherein the vitamin B6 component and the serotonin re-uptake inhibitor are present in a ratio of between 0.01 and 1 mmole/mg, preferably in a ratio of between 0.05 and 0.5 mmole/mg.

10

05.09.21

ABSTRACT

(95)

The present invention is concerned with a method of treating neurological and/or mental disorders that are associated with, or pathogenetically related to deficient serotonin neurotransmission. More particularly the invention relates to a method of increasing cerebral intercellular serotonin levels and/or of preventing decreases in such serotonin levels, said method comprising the administration of serotonin re-uptake inhibitor in a daily amount of between 0.4 and 80 mg and vitamin B6 component in a daily amount of between 0.01 and 10 mmoles, wherein the method does not include the application to the brain of an AC pulsed magnetic field of at least 7.5 picotesia flux density for at least 15 minutes.

The invention also encompasses a pharmaceutical formulation for oral, rectal, buccal or transdermal administration that provides for sustained release of the active agents from the dosage form and a pharmaceutical formulation containing vitamin B6 component and serotonin re-uptake inhibitor a ratio of between 0.01 and 1 mmole/mg.